

Indirect reach of heavy MSSM Higgs bosons by precision measurements at future lepton colliders

Mitsuru Kakizaki,^{1,*} Shinya Kanemura,^{1,†} Mariko Kikuchi,^{1,‡}

Toshinori Matsui,^{1,§} and Hiroshi Yokoya^{1,¶}

¹ *Department of Physics, University of Toyama, Toyama 930-8555, Japan*

(Dated: September 14, 2015)

Abstract

In the Minimal Supersymmetric Standard Model (MSSM), the bottom Yukawa coupling of the Higgs boson can considerably deviate from its Standard Model prediction due to non-decoupling effects. We point out that the ratio of the Higgs boson decay branching fraction to a bottom quark pair and that to a W -boson pair from the same production channel is particularly sensitive to large additional MSSM Higgs boson mass regions at future electron-positron colliders. Based on this precision measurement, we explicitly show the indirect discovery reach of the additional Higgs bosons according to planned programs of the International Linear Collider.

*Electronic address: kakizaki@sci.u-toyama.ac.jp

†Electronic address: kanemu@sci.u-toyama.ac.jp

‡Electronic address: kikuchi@jodo.sci.u-toyama.ac.jp

§Electronic address: matsui@jodo.sci.u-toyama.ac.jp

¶Electronic address: hyokoya@sci.u-toyama.ac.jp; Present address: Theory Center, KEK

In July 2012, discovery of a new particle with a mass of 125 GeV was announced by the ATLAS and CMS collaborations at the CERN Large Hadron Collider (LHC) [1]. Properties of the discovered particle have turned out to be consistent with those of the Higgs boson of the Standard Model (SM) within the error. The SM has been experimentally confirmed as a low-energy effective theory that consistently describes every phenomenon at energy scales below $\mathcal{O}(100)$ GeV. Although the SM is very successful, several phenomena which cannot be explained in the framework of the SM have been known, such as neutrino oscillations, the abundance of dark matter and baryon asymmetry of the Universe. In order to account for such strong evidence we need to go beyond the SM.

It should be emphasized that many new physics models which can explain the new phenomena mentioned above demand extension of the Higgs sector. Given such a non-minimal Higgs sector, a SM-like Higgs boson with a mass of 125 GeV is not exactly the SM one, and has different characteristic properties from their SM predictions. Therefore, it is plausible that hints of new physics are obtained by investigating properties of the discovered Higgs boson. In this viewpoint, the discovered Higgs boson is a window to new physics.

Indeed, one of the best strategies for exploring new physics is to investigate its effects that appear in the couplings of the discovered Higgs boson to SM particles. Since the magnitudes of these Higgs boson couplings depend strongly on masses and couplings of particles yet to be discovered, fingerprinting of the Higgs boson couplings is useful in determining the energy scale of new physics and further in distinguishing new physics models behind. At the LHC, deviations of the Higgs boson couplings from their SM predictions are currently constrained typically with 10% [2]. At future electron-positron colliders such as the International Linear Collider (ILC) [3], the Compact Linear Collider (CLIC) [4] and the Future Circular Collider of electrons and positrons (FCC-ee) [5], accuracies for measurements of various Higgs boson couplings will be significantly improved to a percent level or better [6–8]. Loop corrections from new particles also give characteristic deviation patterns in these couplings. Precision measurements at future collider experiments can differentiate such small deviations. Therefore, it is crucial to perform fingerprinting of the deviation pattern of the Higgs boson couplings based on precise computations including radiative corrections in beyond-the-SM (BSM) models for estimating the scale of new physics indirectly.

Among ever proposed BSM models, supersymmetric (SUSY) extension of the SM is one of the excellent paradigms [9]. Introducing superparticles whose spins are different from the

SM counterparts by one half, an unnatural cancellation between the bare mass squared of the Higgs boson and its quadratic ultraviolet divergences is avoided. By imposing the R -parity, the lightest SUSY particle (LSP) is stabilized and thus can be an excellent candidate for dark matter. The $SU(3)_C$, $SU(2)_L$ and $U(1)_Y$ gauge coupling constants can be naturally unified at a high energy scale, inspiring many ideas about supersymmetric grand unified theories. One of the intriguing features is that the Higgs sector of SUSY models must be extended to cancel the gauge anomaly. Even in the Minimal Supersymmetric Standard Model (MSSM), two Higgs doublets H_1 , H_2 with opposite hypercharges are required. As a consequence, the coupling strengths of the discovered Higgs boson with a mass of 125 GeV to SM particles deviate from their SM predictions, and have the same structure as in the type-II two Higgs doublet model at the tree level. At the loop level, however, superparticles radiatively affect the Higgs boson coupling strengths, giving rise to characteristic deviations patterns different from other models. In particular, it has been known that significant deviations of the bottom and tau Yukawa couplings induced by non-holomorphic radiative corrections do not decouple even in the large superparticle mass limit [10–19]. Due to the non-decoupling effects, the indirect reach of the additional Higgs boson mass scale through precision measurements of the bottom/tau Yukawa coupling can be much higher than the direct reach at the LHC and its luminosity-upgraded version.

In this letter, we reanalyze the deviation of the Higgs boson couplings enhanced by the loop-induced non-decoupling effects from a difference perspective. In sharp contrast to recent relevant works [18, 19], we utilize the ratio of $\text{Br}(h \rightarrow b\bar{b})$ to $\text{Br}(h \rightarrow WW)$ from the same production channel $e^+e^- \rightarrow \nu\bar{\nu}h$ as a key observable. Since this combination cancels out the uncertainties related to the Higgs boson production cross section and total decay rate, the indirect additional Higgs boson reach obtained through this method is more expanded than using single coupling accuracies. Since the cross section times branching ratios of these decay modes are particularly accurately measurable, our quick method is not inferior to global analysis. As an example, we assume expected accuracies at the ILC [8], whose running programs are under discussion. Our goal is to explicitly show the indirect discovery reach of the additional MSSM Higgs bosons through the measurement of the ratio $\text{Br}(h \rightarrow b\bar{b})/\text{Br}(h \rightarrow WW)$ at several planned ILC programs. Such an analysis plays an important role in properly assessing the capability of future electron-positron colliders for exploring new physics effects.

We start briefly reviewing the MSSM Higgs sector, which consists of two Higgs doublet scalars. The electroweak symmetry is broken when the neutral components of H_1 and H_2 develop vacuum expectation values (VEVs). Down-type quarks and charged leptons (up-type quarks) acquire their masses from the VEV $\langle H_1^0 \rangle = v_1/\sqrt{2}$ ($\langle H_2^0 \rangle = v_2/\sqrt{2}$). The ratio $\tan\beta = v_2/v_1$ is an important quantity on which many observables depend, and the sum of the squares is given by $v^2 = v_1^2 + v_2^2 = (246 \text{ GeV})^2$. After the electroweak symmetry breaking, there are two CP -even Higgs bosons h and H , a CP -odd Higgs boson A and charged Higgs bosons $H^\pm$ as physical particles. The lighter CP -even Higgs boson h is identified as the discovered SM-like Higgs boson at the LHC. At the tree level, its predicted mass is smaller than the mass of the Z -boson as the Higgs self-interactions are given by $SU(2)_L$ and $U(1)_Y$ gauge couplings. However, the radiative correction significantly lifts the mass of the Higgs boson and can account for its observed mass of 125 GeV although superparticles are required to have a few TeV masses [20]. The mixing angle α between the two CP -even Higgs bosons is also affected by the radiative correction. For detailed discussion about the radiative corrections to the masses of the CP -even Higgs bosons and its mixing angle, see, for example, Ref. [21]. By virtue of supersymmetry, H , A and $H^\pm$ are degenerate in mass in the large mass limit. We consider the CP -odd Higgs boson mass m_A as a measure of the mass scale of these heavy MSSM Higgs bosons.

Let us discuss non-decoupling effects of radiative corrections to Yukawa couplings in the MSSM. In this letter we focus on the bottom Yukawa coupling because it receives large radiative corrections from the strong coupling and the top Yukawa coupling as well as future linear colliders are capable of measuring the $h \rightarrow b\bar{b}$ decay rate with considerable accuracy. In SUSY models, the holomorphy restricts possible form of the superpotential. As a consequence, the bottom quark does not couple to H_2 at the tree level. However, effects of SUSY breaking induce a non-holomorphic bottom Yukawa coupling $\Delta\lambda_b$ radiatively in addition to holomorphic radiative corrections $\delta\lambda_b$ [14]:

$$-\mathcal{L}_b = (\lambda_b + \delta\lambda_b)\bar{b}_R H_1 Q_L + \Delta\lambda_b \bar{b}_R Q_L H_2^*, \quad (1)$$

where b_R and Q_L denote the right-handed bottom quark and the third generation quark doublet, respectively. Notice that such non-holomorphic couplings are absent in the type-II two-Higgs-doublet model due to the hypothetical Z_2 symmetry. The mass of the bottom

quark is also shifted by the radiative corrections as

$$m_b = \frac{\lambda_b v}{\sqrt{2}} \cos \beta (1 + \Delta_b), \quad (2)$$

where

$$\Delta_b \equiv \frac{\delta \lambda_b}{\lambda_b} + \frac{\Delta \lambda_b}{\lambda_b} \tan \beta. \quad (3)$$

The coupling of the SM-like Higgs boson to the bottom quark is obtained as

$$g_{hb\bar{b}} = \frac{gm_b}{2m_W} \frac{\sin \alpha}{\cos \beta} \left[1 + \frac{1}{1 + \Delta_b} \left(\frac{\delta \lambda_b}{\lambda_b} - \Delta_b \right) (1 + \cot \alpha \cot \beta) \right]. \quad (4)$$

Dominant contributions to the non-holomorphic coupling stem from sbottom-gluino and stop-chargino loop diagrams and are given by

$$\Delta_b \simeq \left(\frac{2\alpha_s}{3\pi} \frac{\mu M_3}{m_{\text{SUSY}}^2} + \frac{\lambda_t^2}{16\pi^2} \frac{\mu A_t}{m_{\text{SUSY}}^2} \right) \tan \beta, \quad (5)$$

where μ , M_3 and A_t are the higgsino mass, the gluino mass and the trilinear stop A -parameter, respectively. The typical mass scale of the superparticles in the loop diagrams is collectively denoted by m_{SUSY} . Notice that there are two mass scales characterizing effects of new physics in the MSSM. One is the mass scale of superparticles m_{SUSY} , and the other is the mass of the CP -odd Higgs boson m_A . In the limit where superparticles are heavy $\mu \sim M_3 \sim A_t \sim m_{\text{SUSY}} \gg m_Z$ with fixed m_A , the bottom Yukawa coupling is approximately given by

$$g_{hb\bar{b}} \simeq \frac{gm_b}{2m_W} \frac{\sin \alpha}{\cos \beta} [1 - \Delta_b (1 + \cot \alpha \cot \beta)]. \quad (6)$$

Therefore, SUSY loop corrections do not decouple for small m_A , and are enhanced for large $\tan \beta$. On the other hand, in the limit where the mass of the CP -odd Higgs boson is large, we obtain

$$1 + \cot \alpha \cot \beta = -\frac{2m_Z^2}{m_A^2} \cos 2\beta + \mathcal{O}\left(\frac{m_Z^4}{m_A^4}\right), \quad (7)$$

and thus SUSY radiative corrections decouple. It should be noticed that two-Higgs-doublet models can also produce non-decoupling effect on the Higgs boson couplings, which originates from extra Higgs boson loop diagrams [22–26].

We here describe the method for our numerical analysis. We utilize the public code FeynHiggs2.10.3 to compute mass and mixing parameters as well as decay branching fractions in the MSSM Higgs sector [27]¹. The masses of the CP -even Higgs bosons and the mixing angle are computed at the two-loop level in the on-shell scheme. Branching ratios of the SM-like Higgs boson are computed at the one-loop level including QED and QCD corrections. We perform a random scan of MSSM parameters avoiding tachyonic scalar masses. To circumvent large CP -violating processes, we constrain our analysis to real MSSM parameters. For simplicity, we assume that soft SUSY breaking A -parameters are given by the product of the corresponding Yukawa coupling constant λ_f and a common A_0 value as $A_f = \lambda_f A_0$. We consider the mass range of electroweak (colored) superparticles up to around 1000 GeV (4000 GeV) to cover less tuned parameter sets which evade superparticle searches at the future LHC experiment. Since the non-decoupling effects depend on ratios of superparticle masses, our results about the Higgs boson couplings are not substantially changed even for heavier superparticles. If $|A_t|$ is significantly larger than the stop masses, there appear a deeper charge/color breaking vacuum than our electroweak vacuum. To avoid complexity of the discussion of the meta-stability bound, we constrain $|A_0|$ to be smaller than three times the geometric average of the soft stop masses. The scan bounds of the MSSM parameters are listed in Tab. I. Since radiative corrections from the first two generation sleptons and squarks to the Higgs sector are small, we set the soft masses of these sleptons and squarks to 1000 GeV and 4000 GeV, respectively, instead of performing a random scan. This degeneracy also suppresses unwanted flavor chaining neutral currents and lepton flavor violating processes adequately. Although the measured mass of the Higgs boson is now fixed to $m_h = 125$ GeV with the error less than 1 GeV [28], we consider $122 \text{ GeV} < m_h < 128 \text{ GeV}$ as an allowed region for its mass. This is a judicious choice to take uncertainties from renormalization scheme dependence [29] and higher order contributions to the mass of the Higgs boson into account [30]. From the cosmological viewpoint, it is assumed that the lightest neutralino is the LSP and a dark matter candidate. However, the relic density constraint is relaxed in

¹ Changes in the recently updated version FeynHiggs2.11.0 are not relevant to our analysis.

TABLE I: Scan bounds on MSSM parameters.

Parameter	Scan bounds
m_A	[200 GeV, 3000 GeV]
$\tan \beta$	[1, 60]
$\tilde{m}_{L_3, E_3}$	[100 GeV, 1000 GeV]
$\tilde{m}_{Q_3, U_3, D_3}$	[500 GeV, 4000 GeV]
$ A_0 $	[0 GeV, $3(\tilde{m}_{Q_3}\tilde{m}_{U_3})^{1/2}$]
$ \mu $	[100 GeV, 1000 GeV]
$ M_1 $	[100 GeV, 1000 GeV]
$ M_2 $	[100 GeV, 1000 GeV]
M_3	[1400 GeV, 4000 GeV]

TABLE II: Constraints on superparticle masses.

Observables	Constraints
m_h	[122 GeV, 128 GeV]
$m_{\tilde{q}}$	> 1400 GeV
$m_{\tilde{\chi}_1^0}$	> 90 GeV
$m_{\tilde{\chi}_1^\pm}$	> 400 GeV
$m_{\tilde{t}_1}$	> 640 GeV
$m_{\tilde{b}_1}$	> 620 GeV
$m_{\tilde{\tau}_1}$	> 90 GeV

order to allow for any non-standard cosmological scenario. As for the other experimental constraints, we impose superparticle mass bounds obtained at the LEP [31] and the LHC [32]. The mass bounds we employ are summarized in Tab. II. The flavor violating decays $B_s \rightarrow \mu^+ \mu^-$ and $b \rightarrow s \gamma$ mediated by the additional Higgs bosons could also give constraints on the SUSY parameter space, as studied in [19]. However, such processes can be canceled by contributions from squark sector parameters without changing our results. Therefore, we neglect these flavor constraints in our analysis. The deviation of the anomalous magnetic moment of the muon from its SM prediction can be explained by adjusting the masses of the second generation sleptons, which is beyond the scope of this letter. Since our analysis is confined to low-energy SUSY scenarios, we require λ_b to be perturbative at the electroweak scale.

Our goal is to show to what extent large m_A scenarios with moderate SUSY parame-

TABLE III: Expected accuracy for R at planned stages of the ILC [8].

CM energy and luminosity	95%CL	5σ
$\sqrt{s} = 500 \text{ GeV}, L = 500 \text{ fb}^{-1}$	0.050	0.125
$\sqrt{s} = 500 \text{ GeV}, L = 1600 \text{ fb}^{-1}$	0.027	0.068
$\sqrt{s} = 1 \text{ TeV}, L = 2500 \text{ fb}^{-1}$	0.021	0.052

ter choices can be surveyed through the precision measurements of the cross section times branching ratio measurements of the $h \rightarrow b\bar{b}$ and $h \rightarrow WW$ modes which follow the production $e^+e^- \rightarrow \nu\bar{\nu}h$. To this end, we introduce the following double ratio of the branching fractions of $h \rightarrow b\bar{b}$ and that of $h \rightarrow WW$ modes:

$$R = \frac{\text{Br}(h \rightarrow WW)}{\text{Br}(h \rightarrow b\bar{b})} \bigg|_{\text{MSSM}} \bigg/ \frac{\text{Br}(h \rightarrow WW)}{\text{Br}(h \rightarrow b\bar{b})} \bigg|_{\text{SM}}. \quad (8)$$

Its tree level expression is given by

$$R_{\text{tree}} = \frac{1}{[1 - \tan \beta \cot(\beta - \alpha)]^2}, \quad (9)$$

where $\cot(\beta - \alpha)$ is given as a function of m_A and m_Z . As is evident, any deviation from the SM prediction $R = 1$ exhibits the existence of some new physics beyond the SM. By taking the ratio of the cross section times branching ratios for $b\bar{b}$ and WW modes in the same production mechanism, uncertainties that originate from the production cross section and the total decay rate of the Higgs boson are canceled out. Consequently, constraining the model parameter space using the accuracy of the ratio of the decay branching fraction R is more powerful than using the converted accuracy of the $hb\bar{b}$ coupling. The expected accuracies for cross section times branching ratio measurements of the SM-like Higgs boson through the $\nu\bar{\nu}h$ mode at several ILC stages with the polarization options $(P_{e^-}, P_{e^+}) = (-0.8, +0.3)$ and $(P_{e^-}, P_{e^+}) = (-0.8, +0.2)$ are listed in Tab. 5.4 and 5.5 of Ref. [8]. Sensitivities for R are computed based on these tables and summarized in Tab. III. For example, for $\sqrt{s} = 500 \text{ GeV}$ and $L = 500 \text{ fb}^{-1}$, the accuracy of the ratio at 95% CL is estimated as $\sqrt{(0.7\%)^2 + (2.4\%)^2} \times 2 = 5\%$, while the accuracy of $\text{Br}(h \rightarrow b\bar{b})$ from the coupling measurement itself is at best $\simeq 6.4\%$ ignoring systematic uncertainties (see Table 6.1 in Ref.[8]).

In Fig. 1, the indirect 5σ discovery reach of the MSSM heavy Higgs bosons through the measurement of R at the ILC with the center-of-mass (CM) energy $\sqrt{s} = 500$ GeV and the luminosity $L = 500 \text{ fb}^{-1}$ is shown in the $(m_A, \tan\beta)$ plane. MSSM scenarios which predict more than 5σ deviation of R are plotted with the red points. The blue line is the corresponding tree level prediction based on R_{tree} . The indirect reach of the MSSM heavy Higgs boson mass can be significantly extended due to the non-decoupling property of the non-holomorphic radiative corrections, compared to the tree level results. When the relative sign of μ and M_3 (A_t) is minus, the deviation of R from unity is strongly enhanced and the correction is proportional to $\tan\beta$ (see Eqs.(5) and (6)). The solid (black) line shows the 95% exclusion limit obtained at the LHC Run-I through direct $H/A \rightarrow \tau^+\tau^-$ searches [33]. At the LHC Run-II with the luminosity of $L = 10 \text{ fb}^{-1}$ accumulated, it is expected to exclude up to $m_A = 1000$ GeV for $\tan\beta = 10$ in view of the increase of the production cross section [34]. The existence of the heavy Higgs boson can be searched also through the $H/A \rightarrow \mu^+\mu^-$ decay modes at the LHC [35]. Its expected 5σ discovery reach with $L = 3000 \text{ fb}^{-1}$ almost overlaps with the 95% excluded region explored by the τ channel.

The 5σ heavy Higgs mass reach expected at the stage with $\sqrt{s} = 500$ GeV and $L = 1600 \text{ fb}^{-1}$ and at the stage with $\sqrt{s} = 1$ TeV and $L = 2500 \text{ fb}^{-1}$ are shown in Fig. 2 and Fig. 3, respectively. These figures show that luminosity-upgraded ILC programs are very powerful in measuring the Higgs boson branching fractions and thus sensitive to higher m_A regions [8].

Before closing, we comment on the parametric uncertainties from the SM parameters [36]. Theoretical calculation on $\text{Br}(h \rightarrow b\bar{b})$ suffers from uncertainties on m_b and α_s , and the uncertainties on the calculation limit the reach of MSSM parameters by experimental measurements [37]. Although current errors on these parameters [38] lead to larger uncertainties than the expected uncertainties at the future ILC measurements, it is anticipated that future developments on higher order calculation and lattice calculation overcome this situation until the experimental programs take place [39].

In this letter, we have presented to what extent the heavy additional MSSM Higgs bosons can be indirectly accessible by using precision measurements of the cross section times branching ratios for the discovered Higgs boson at the future ILC experiments. We have employed the ratio of the Higgs boson decay branching fractions $\text{Br}(h \rightarrow b\bar{b})/\text{Br}(h \rightarrow WW)$ as a simple but useful observable for indirectly probing the existence of the heavy Higgs




FIG. 1: Indirect 5σ discovery reach of the heavy Higgs bosons through the measurement of R at the ILC stage with $\sqrt{s} = 500 \text{ GeV}$ and $L = 500 \text{ fb}^{-1}$ in the $(m_A, \tan \beta)$ plane (red points). For comparison, the indirect 5σ reach based on the tree-level results is delineated with blue line. The 95% exclusion limit obtained at the LHC Run-I through $H/A \rightarrow \tau^+\tau^-$ searches is delineated with black line.

bosons because this ratio is obtained by using the cross section times branching fractions, which cancel out the uncertainties of the production cross section and total decay width of the Higgs boson. We have computed the ratio R , which can substantially deviate from unity even for large m_A regions due to large non-decoupling effects of superparticles, and have exhibited the indirect 5σ discovery potential of the additional MSSM Higgs bosons at several planned stages of the future ILC experiments. Our analysis shows that the ILC, in particular with high luminosity, cover many large m_A scenarios in the MSSM.

Last but not least, fingerprinting of Higgs boson couplings at the loop level including our analysis improves capability of exploring and distinguishing the MSSM from other models of the electroweak symmetry breaking, and should be encouraged.

Acknowledgments

The authors would like to thank Hiroyuki Taniguchi for early stage collaboration. The work of M. Kakizaki was supported in part by Grant-in-Aid for Scientific Research, No. 26104702. The work of S.K. was supported in part by Grant-in-Aid for Scientific Research,




FIG. 2: Indirect 5σ discovery reach of the heavy Higgs bosons through the measurement of R at the luminosity-upgraded ILC stage with $\sqrt{s} = 500 \text{ GeV}$ and $L = 1600 \text{ fb}^{-1}$ in the $(m_A, \tan\beta)$ plane (red points). The others are same as in Fig.1.

Nos. 24340046, 23104006 and 30181308. The work of M. Kikuchi was supported in part by Japan Society for the Promotion of Science, No. 25·10031. The work of T.M. was supported in part by the Sasakawa Scientific Research Grant from The Japan Science Society.

-
- [1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012); S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012).
 - [2] V. Khachatryan *et al.* [CMS Collaboration], arXiv:1412.8662 [hep-ex]; ATLAS Collaboration, ATLAS-CONF-2015-007.
 - [3] J. Brau, (Ed.) *et al.* [ILC Collaboration], arXiv:0712.1950 [physics.acc-ph]; G. Aarons *et al.* [ILC Collaboration], arXiv:0709.1893 [hep-ph]; N. Phinney, N. Toge and N. Walker, arXiv:0712.2361 [physics.acc-ph]; T. Behnke, (Ed.) *et al.* [ILC Collaboration], arXiv:0712.2356 [physics.ins-det]; T. Behnke *et al.*, arXiv:1306.6329 [physics.ins-det]; H. Baer, *et al.* "Physics at the International Linear Collider", *Physics Chapter of the ILC Detailed Baseline Design Report*: <http://lcsim.org/papers/DBDPhysics.pdf>.




FIG. 3: Indirect 5σ discovery reach of the heavy Higgs bosons through the measurement of R at the luminosity-upgraded ILC stage with $\sqrt{s} = 1 \text{ TeV}$ and $L = 2500 \text{ fb}^{-1}$ in the $(m_A, \tan \beta)$ plane (red points). The others are same as in Fig.1.

- [4] E. Accomando *et al.* [CLIC Physics Working Group Collaboration], hep-ph/0412251; L. Linssen, A. Miyamoto, M. Stanitzki and H. Weerts, arXiv:1202.5940 [physics.ins-det].
- [5] M. Bicer *et al.* [TLEP Design Study Working Group Collaboration], JHEP **1401** (2014) 164.
- [6] M. E. Peskin, arXiv:1207.2516 [hep-ph].
- [7] H. Baer, T. Barklow, K. Fujii, Y. Gao, A. Hoang, S. Kanemura, J. List and H. E. Logan *et al.*, arXiv:1306.6352 [hep-ph].
- [8] D. M. Asner, T. Barklow, C. Calancha, K. Fujii, N. Graf, H. E. Haber, A. Ishikawa and S. Kanemura *et al.*, arXiv:1310.0763 [hep-ph].
- [9] For reviews of SUSY phenomenology, see for example, S. P. Martin, Adv. Ser. Direct. High Energy Phys. **21**, 1 (2010) [hep-ph/9709356]; M. Drees, R. Godbole and P. Roy, Hackensack, USA: World Scientific (2004) 555 p; H. Baer and X. Tata, Cambridge, UK: Univ. Pr. (2006) 537 p.
- [10] L. J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. D **50**, 7048 (1994) [hep-ph/9306309, hep-ph/9306309].
- [11] R. Hempfling, Phys. Rev. D **49**, 6168 (1994).

- [12] M. Carena, M. Olechowski, S. Pokorski and C. E. M. Wagner, Nucl. Phys. B **426**, 269 (1994) [hep-ph/9402253].
- [13] M. Carena, S. Mrenna and C. E. M. Wagner, Phys. Rev. D **60**, 075010 (1999) [hep-ph/9808312].
- [14] K. S. Babu and C. F. Kolda, Phys. Lett. B **451**, 77 (1999) [hep-ph/9811308].
- [15] H. Eberl, K. Hidaka, S. Kraml, W. Majerotto and Y. Yamada, Phys. Rev. D **62** (2000) 055006 [hep-ph/9912463].
- [16] M. Carena, D. Garcia, U. Nierste and C. E. M. Wagner, Nucl. Phys. B **577**, 88 (2000) [hep-ph/9912516].
- [17] H. E. Haber and J. D. Mason, Phys. Rev. D **77**, 115011 (2008) [arXiv:0711.2890 [hep-ph]].
- [18] M. Cahill-Rowley, J. Hewett, A. Ismail and T. Rizzo, Phys. Rev. D **90**, no. 9, 095017 (2014) [arXiv:1407.7021 [hep-ph]].
- [19] M. Endo, T. Moroi and M. M. Nojiri, arXiv:1502.03959 [hep-ph].
- [20] Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991); J. R. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. B **257**, 83 (1991); H. E. Haber and R. Hempfling, Phys. Rev. Lett. **66**, 1815 (1991).
- [21] A. Djouadi, Phys. Rept. **459**, 1 (2008) [hep-ph/0503173].
- [22] S. Kanemura, S. Kiyoura, Y. Okada, E. Senaha and C. P. Yuan, Phys. Lett. B **558**, 157 (2003) [hep-ph/0211308]; S. Kanemura, Y. Okada, E. Senaha and C.-P. Yuan, Phys. Rev. D **70**, 115002 (2004) [hep-ph/0408364].
- [23] A. Arhrib, M. Capdequi Peyranere, W. Hollik and S. Penaranda, Phys. Lett. B **579**, 361 (2004) [hep-ph/0307391].
- [24] S. Kanemura, M. Kikuchi and K. Yagyu, Phys. Lett. B **731**, 27 (2014) [arXiv:1401.0515 [hep-ph]]; Nucl. Phys. B **896**, 80 (2015) [arXiv:1502.07716 [hep-ph]].
- [25] P. Osland, P. N. Pandita and L. Selbuz, Phys. Rev. D **78**, 015003 (2008) [arXiv:0802.0060 [hep-ph]].
- [26] L. Wu, J. M. Yang, C. P. Yuan and M. Zhang, arXiv:1504.06932 [hep-ph].
- [27] S. Heinemeyer, W. Hollik and G. Weiglein, Comput. Phys. Commun. **124**, 76 (2000) [hep-ph/9812320]; S. Heinemeyer, W. Hollik and G. Weiglein, Eur. Phys. J. C **9**, 343 (1999) [hep-ph/9812472]; G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, Eur. Phys. J. C **28**, 133 (2003) [hep-ph/0212020]; M. Frank, T. Hahn, S. Heinemeyer, W. Hollik,

- H. Rzehak and G. Weiglein, JHEP **0702**, 047 (2007) [hep-ph/0611326]; T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak and G. Weiglein, Phys. Rev. Lett. **112**, no. 14, 141801 (2014) [arXiv:1312.4937 [hep-ph]].
- [28] G. Aad *et al.* [ATLAS and CMS Collaborations], arXiv:1503.07589 [hep-ex].
- [29] For recent analysis, see, for example, L. J. Hall, D. Pinner and J. T. Ruderman, JHEP **1204**, 131 (2012); P. Draper, P. Meade, M. Reece and D. Shih, Phys. Rev. D **85**, 095007 (2012); S. Borowka, T. Hahn, S. Heinemeyer, G. Heinrich and W. Hollik, Eur. Phys. J. C **74**, no. 8, 2994 (2014) [arXiv:1404.7074 [hep-ph]]; G. Degrossi, S. Di Vita and P. Slavich, Eur. Phys. J. C **75**, no. 2, 61 (2015) [arXiv:1410.3432 [hep-ph]]; S. Borowka, T. Hahn, S. Heinemeyer, G. Heinrich and W. Hollik, arXiv:1505.03133 [hep-ph].
- [30] R. V. Harlander, P. Kant, L. Mihaila and M. Steinhauser, Phys. Rev. Lett. **100**, 191602 (2008) [Phys. Rev. Lett. **101**, 039901 (2008)] [arXiv:0803.0672 [hep-ph]]; JHEP **1008**, 104 (2010) [arXiv:1005.5709 [hep-ph]]; J. L. Feng, P. Kant, S. Profumo and D. Sanford, Phys. Rev. Lett. **111** (2013) 131802 [arXiv:1306.2318 [hep-ph]].
- [31] LEPSUSYWG, ALEPH, DELPHI, L3 and OPAL experiments, notes LEPSUSYWG/04-01.1, LEPSUSYWG/01-03.1, and LEPSUSYWG/02-04.1.
- [32] F. Würthwein, plenary talk at ICHEP 2014, Valencia, Spain, 2-9 July 2014.
- [33] V. Khachatryan *et al.* [CMS Collaboration], JHEP **1410**, 160 (2014) [arXiv:1408.3316 [hep-ex]]; G. Aad *et al.* [ATLAS Collaboration], JHEP **1411**, 056 (2014) [arXiv:1409.6064 [hep-ex]].
- [34] M. Morinaga, private communication.
- [35] ATLAS Collaboration, ATLAS-PHYS-PUB-2013-016.
- [36] A. Droll and H. E. Logan, Phys. Rev. D **76** (2007) 015001 [hep-ph/0612317]; A. Denner, S. Heinemeyer, I. Puljak, D. Rebuszi and M. Spira, Eur. Phys. J. C **71** (2011) 1753 [arXiv:1107.5909 [hep-ph]]; L. G. Almeida, S. J. Lee, S. Pokorski and J. D. Wells, Phys. Rev. D **89** (2014) 3, 033006 [arXiv:1311.6721 [hep-ph]].
- [37] K. Desch, E. Gross, S. Heinemeyer, G. Weiglein and L. Zivkovic, JHEP **0409** (2004) 062 [hep-ph/0406322].
- [38] S. Eidelman *et al.* [Particle Data Group Collaboration], Phys. Lett. B **592** (2004) 1.
- [39] G. P. Lepage, P. B. Mackenzie and M. E. Peskin, arXiv:1404.0319 [hep-ph].